



University of Groningen

## Reply: Hering's Law of the Frontal Facial Branch

Gülbitti, Haydar Aslan; van der Lei, Berend

*Published in:*

Plastic and Reconstructive Surgery

*DOI:*

[10.1097/PRS.00000000000005974](https://doi.org/10.1097/PRS.00000000000005974)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*

Final author's version (accepted by publisher, after peer review)

*Publication date:*

2019

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Gülbitti, H. A., & van der Lei, B. (2019). Reply: Hering's Law of the Frontal Facial Branch. *Plastic and Reconstructive Surgery*, 144(3), 519E-520E. <https://doi.org/10.1097/PRS.00000000000005974>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

**Re: Reply to Hering's Law of the Frontal Facial Branch**

Haydar Aslan Gülbitti<sup>1</sup>, M.D. and Berend van der Lei<sup>1, 2</sup>, M.D., Ph. D.

1. Department of Plastic Surgery, University of Groningen, University Medical Centre  
Groningen, Groningen, The Netherlands

2. Bey Bergman Clinics, Hilversum, The Netherlands

**Running Title:** Re: Reply to Hering's Law of the Frontal Facial Branch

(Key Words: Hering's Law, Facial Palsy)

There has been no financial support and there are no financial interests.

**Address for correspondence:** Prof. Dr. B. van der Lei, University Medical Center

Groningen, PO box 30.001, 9700 RB Groningen, fax. (+31)(0)50-3613043, tel. (+31)(0)50-3613531.

Sir,

We thank prof. Procianoy for his insightful comments on our short article concerning Hering's law of the frontal facial branch<sup>1</sup>. We fully agree that there is a lack of scientific evidence in literature to prove the pathophysiology of our findings. It is indeed hard to prove that contralateral elevation of the eyebrow is induced by either voluntary or involuntary contraction on an electromyographic level. However, according to our opinion, elevation of the contralateral brow due to voluntary contraction after ipsilateral paralysis is very much unlikely; in the case of voluntary contraction, the patient will attempt to elevate the affected side of the face, not the contralateral side.

Although it is understandable that prof. Procianoy uses the manuscript of Beaulieu, Andre and Mancini for his argumentation, an in depth reading of this respective article would quickly lead to the conclusion that their presented results do not reflect to ours<sup>2</sup>. This is due to the fact that their study population consisted of healthy volunteers, whereas our study population consisted of facial paralyzed patients. Their study showed that the eyebrow position does not change when the visual field was deprived via black contact lenses and when an external weight was placed on the eyelid the patient attempted to raise the both eyelid and eyebrow to obtain the same eyesight. As expected, eyebrow position will not change in healthy volunteers when the sight is covered by an object (such as a lens or glasses), however, the ipsilateral eyebrow position will (voluntary) elevate in attempt to raise against external weight, as explained before. Patients with facial paralysis also attempt to raise the affected eyebrow, like in healthy volunteers, but this is off course impossible due to ipsilateral impairment of the facial nerve. Therefore, in case of facial nerve palsy, we see that the contralateral eyebrow continuously is elevated and indicates that Hering's law also applies to the frontal branch of the facial nerve. Moreover, blocking the eyesight on the paralyzed side results in lowering of the eyebrow of the contralateral side.

There is no scientific evidence that muscular hypertrophy occurs in facial paralyzed patients, as mentioned by prof. Procianoy. Choe et al. and Choi et al., however, only have suggested this by writing: “we treated 34 patients who had a drooping face on one side after facial paralysis with thread lifting and botulinum injections to control contralateral hypertrophy and ipsilateral synkinesis” respectively “the non-paralysed side, which acts chronically against the weak antagonism of the contralateral muscles, usually presents with facial muscular hypertrophy, wrinkles, furrows and deviation of the mouth. Facial changes associated with facial sequelae can be attributed to ‘unbalanced’ muscular activity, ‘unbalanced’ muscular hypertrophy and ‘unbalanced’ pattern of facial expression”<sup>3,4</sup>.

Further studies, including electromyographic ones, for sure are warranted to further elucidate Hering’s law of the frontal facial branch and that of perhaps the other facial branches.

## References

1. Gülbitti HA, van der Lei B. Hering's Law of the Frontal Facial Branch. *Plast Reconstr Surg*. 2018 Dec;142(6):991e-992e.
2. Frontalis Muscle Contraction and the Role of Visual Deprivation and Eyelid Proprioception. Beaulieu R, Andre K, Mancini R. *Ophthalmic Plast Reconstr Surg*. 2018 Nov/Dec;34(6):552-556.
3. Choe WJ, Kim HD, Han BH, Kim J. Thread lifting: a minimally invasive surgical technique for long-standing facial paralysis. *HNO*. 2017 Nov;65(11):910-915.
4. Choi KH, Rho SH, Lee JM, Jeon JH, Park SY, Kim J. Botulinum toxin injection of both sides of the face to treat post-paralytic facial synkinesis. *J Plast Reconstr Aesthet Surg*. 2013 Aug;66(8):1058-63.